



ELSEVIER

Resuscitation 57 (2003) 231–235

RESUSCITATION



www.elsevier.com/locate/resuscitation

Therapeutic hypothermia after cardiac arrest. An advisory statement by the Advanced Life Support Task Force of the International Liaison Committee on Resuscitation[☆]

Jerry P. Nolan^{a,*}, Peter T. Morley^b, Terry L. Vanden Hoek^c, Robert W. Hickey^{d,1},
ALS Task Force²

^a *Cochair ILCOR, Department of Anaesthesia and Intensive Care Medicine, Royal United Hospital, Bath BA1 3NG, UK*

^b *Chairman, Advanced Life Support Committee, Australian Resuscitation Council, Intensive Care Unit, Royal Melbourne Hospital, Melbourne, Vic. 3050, Australia*

^c *Member, ACLS Subcommittee, American Heart Association, Assistant Professor of Emergency Medicine, University of Chicago, 5841 South Maryland Ave, MC5068, Chicago, IL 60637, USA*

^d *Chair, Subcommittee on Pediatric Resuscitation, American Heart Association, Associate Professor of Pediatrics, Children's Hospital of Pittsburgh, Division of Pediatric Emergency Medicine, 3705 Fifth Avenue, Pittsburgh, PA 15213, USA*

1. ILCOR recommendations

On the basis of the published evidence to date, the Advanced Life Support (ALS) Task Force of the International Liaison Committee on Resuscitation

(ILCOR) made the following recommendations in October 2002:

- Unconscious adult patients with spontaneous circulation after out-of-hospital cardiac arrest should be cooled to 32–34 °C for 12–24 h when the initial rhythm was ventricular fibrillation (VF).
- Such cooling may also be beneficial for other rhythms or in-hospital cardiac arrest.

[☆] Every effort has been made to avoid any actual or potential conflicts of interest that may arise as a result of an outside relationship or a personal, professional, or business interest of a member of the writing panel. Specifically, all members of the writing group are required to complete and submit a Disclosure Questionnaire showing all such relationships that might be perceived as real or potential conflicts of interest. This statement has been co-published in the 8 July 2003, issue of *Circulation*. This statement was approved by the Advanced Life Support Task Force of the International Liaison Committee on Resuscitation in April 2003.

* Corresponding author.

E-mail address: jerry.nolan@ruh-bath.swest.nhs.uk (J.P. Nolan).

¹ Member of the Pediatric Life Support Task Force.

² Members of the ALS Task Force: W.G.J. Kloeck, Chair (Resuscitation Council of Southern Africa [RCSA]); J. Billi, MD (American Heart Association [AHA]); B.W. Böttiger (European Resuscitation Council [ERC]); P.T. Morley (Australia and New Zealand Council on Resuscitation [ANZCOR]); J.P. Nolan (Cochair, ILCOR); K. Okada (Japanese Resuscitation Council [JRC]); C. Reyes, MD (Latin American Resuscitation Council [CLAR]); M. Shuster, MD, FRCPC (Heart and Stroke Foundation of Canada [HSFC]); P.A. Steen, MD (ERC); M.H. Weil, MD, PhD (AHA); V. Wenzel (ERC). Also additional contributors: P. Carli (ERC), T.L. Vanden Hoek, MD (AHA), D. Atkins, MD (AHA).

2. Introduction

Induction of moderate hypothermia (28–32 °C) before cardiac arrest has been used successfully since the 1950s to protect the brain against the global ischaemia that occurs during some open-heart surgery. Successful use of therapeutic hypothermia after cardiac arrest in humans was also described in the late 1950s [1–3] but was subsequently abandoned because of uncertain benefit and difficulties with its use [4]. Since then, induction of hypothermia after return of spontaneous circulation (ROSC) has been associated with improved functional recovery and reduced cerebral histological deficits in various animal models of cardiac arrest [5–8]. Additional promising preliminary human studies have been completed [9–16]. At the time of publication of the *Guidelines 2000 for Cardiopulmonary Resuscitation and*

Emergency Cardiovascular Care, the evidence was insufficient to recommend use of therapeutic hypothermia after resuscitation from cardiac arrest [17].

3. Clinical studies

In 2002 the results of two prospective randomised trials were published that compared mild hypothermia with normothermia in comatose survivors of out-of-hospital cardiac arrest [18,19]. One study was undertaken in nine centers in five European countries [19]; the other was conducted in four hospitals in Melbourne, Australia [18].

The criteria for entry into these trials were similar: ROSC, patients remaining intubated and ventilated, with persistent coma after out-of-hospital cardiac arrest due to VF. In the European study, the median Glasgow Coma Scale score on hospital admission in both groups was 3 with an interquartile range of 3–5 in the group with normothermia and 3–4 in the group with hypothermia [20]. Ten patients in the European study were resuscitated after in-hospital cardiac arrest (M. Holzer, written communication, October 2002). Additional criteria for entry into the European study were witnessed cardiac arrest, an estimated interval of 5–15 min from patient collapse to first resuscitation attempt by emergency medical services personnel, and an interval of ≤ 60 min from collapse to ROSC. Both studies excluded arrests that were probably of noncardiac aetiology and patients with severe cardiogenic shock.

In the European study, patients randomly assigned to the hypothermia group underwent cooling to a target temperature of 32–34 °C by use of a mattress made specifically for this purpose (with a cover that delivered cold air) and ice packs if necessary. The aim was to reach the target temperature within 4 h of ROSC, maintain it for 24 h, and follow with passive rewarming. In the Australian study, patients were pseudo-randomised (odd vs. even days) to treatment groups that allowed cooling, by application of cold packs to the head and torso, to begin in the field before admission to the hospital. The target temperature of the hypothermia group was 33 versus 37 °C in the control group. Hypothermia was maintained for 12 h after admission to the hospital; active rewarming started at 18 h.

In the European study, 75 of the 136 patients (55%) in the hypothermia group for whom data were available had a favorable neurological outcome (able to live independently and work at least part-time) at 6 months compared with 54 of 137 (39%) in the normothermia group (relative risk [RR] 1.40, 95% CI 1.08–1.81, number needed to treat [NNT] = 6) [19]. At 6 months there were 56 deaths in the 137 participants (41%) in the hypothermia group versus 76 of 138 (55%) in the normothermia group (RR 0.74, 95% CI 0.58–0.95,

NNT = 7). The target temperature could not be achieved in 19 patients in the hypothermia group.

In the Australian study, 21 of 43 patients (49%) treated with hypothermia had good neurological function at discharge (to home or a rehabilitation facility) compared with 9 of 34 (26%) in the normothermia group (RR 1.85, 95% CI 0.97–3.49, NNT = 4) [18]. Mortality at discharge was 22 of 43 (51%) in the hypothermia group and 23 of 34 (68%) in the normothermia group (RR 0.76, 95% CI 0.52–1.10, NNT = 6).

Both of these studies involved a highly selected group of patients, excluding up to 92% of patients with out-of-hospital cardiac arrest initially assessed for eligibility [19]. Those excluded had persistent hypotension (systolic blood pressure < 90 mmHg despite use of inotropes) and causes of coma other than cardiac arrest (e.g. head injury, drug overdose, cerebrovascular accident). Other study limitations were that caregivers could not be blinded to treatment with hypothermia and that following ROSC, the normothermia group had an increase in core temperature of up to > 38 °C, as is often seen after cardiac arrest.

Some adverse events occurred more frequently in the hypothermia groups. The Australian hypothermia group had a lower cardiac index, higher systemic vascular resistance, and more hyperglycaemia than patients in the control group [18]. Although not statistically significant, in the European hypothermia group there were 22% more complications; in particular, there were more cases of pneumonia [number needed to harm (NNH) = 12], bleeding (NNH = 14), and sepsis (NNH = 16) [19].

4. Mechanisms of action

There are several possible mechanisms by which mild hypothermia might improve neurological outcome when used after reperfusion. In the normal brain, hypothermia reduces the cerebral metabolic rate for oxygen (CMRO₂) by 6% for every 1 °C reduction in brain temperature > 28 °C [21]. Some of this effect is due to reduced normal electrical activity [21], however, and after cardiac arrest in dogs, CMRO₂ is not significantly reduced by mild hypothermia [22]. Mild hypothermia is thought to suppress many of the chemical reactions associated with reperfusion injury. These reactions include free radical production, excitatory amino acid release, and calcium shifts, which can in turn lead to mitochondrial damage and apoptosis (programmed cell death) [23–25]. Despite these potential advantages, hypothermia can also produce adverse effects, including arrhythmias, infection, and coagulopathy.

5. Discussion

5.1. Selection of patients

There seems to be good evidence (Level 1 [see Appendix A]) to recommend the use of induced mild hypothermia in comatose survivors of out-of-hospital cardiac arrest caused by VF. Selection criteria for treatment were narrowly defined in the best evidence used and thus should be considered carefully when deciding to treat.

Several specific questions remain unanswered despite the results of these recently published controlled trials, previous clinical studies, and supporting experiments in animals. One controversial issue is whether findings from animal experiments and published clinical studies are enough to extend the use of therapeutic mild hypothermia to patients who remain comatose after cardiac arrest from any rhythm, after in-hospital cardiac arrest, and after cardiac arrest in children.

Any potentially beneficial effects of hypothermia on neuronal recovery must be counterbalanced by the known adverse effects of hypothermia. Although survivors of VF cardiac arrest have the most to gain from therapeutic hypothermia, some level 4 evidence suggests that survivors from out-of-hospital cardiac arrest of other causes may also benefit [9]. Further study is required. Many in-hospital cardiac arrests have non-cardiac causes, and because the use of therapeutic hypothermia has not been studied to a significant extent in this population, its relative risks and benefits are unknown. It is possible, however, that patients who remain comatose after an in-hospital arrest of cardiac aetiology will also benefit from therapeutic hypothermia.

Until further data are available, therapeutic hypothermia should not be used for patients with severe cardiogenic shock or life-threatening arrhythmias, pregnant patients, or patients with primary coagulopathy. Thrombolytic therapy does not preclude the use of hypothermia; patients who received thrombolytic therapy were included in both the European and Australian studies.

5.2. Timing of cooling

Cooling should probably be initiated as soon as possible after ROSC but appears to be successful even if delayed (e.g. 4–6 h). In the European study, the interval between ROSC and attainment of a core temperature of 32–34 °C had an interquartile range of 4–16 h [19].

Further research is needed to determine optimal duration of therapeutic hypothermia, optimum target temperature, and rates of cooling and rewarming.

Animal data suggest that the sooner cooling is initiated after reperfusion from cardiac arrest, the better the outcome, although an impressive therapeutic benefit was seen in clinical studies when cooling was delayed for several hours. The therapeutic benefit may become much greater as better physical and pharmacological techniques to cool patients more rapidly become available. Although supporting data are limited, many critical care clinicians routinely sedate and ventilate the lungs of comatose survivors of cardiac arrest for at least 12–24 h; thus, application of therapeutic hypothermia over this period would be simple. Normothermia should be restored only slowly as rebound hyperthermia is common and should be avoided [14].

5.3. Cooling techniques and monitoring

A variety of cooling techniques have been described, but at this stage, none combines ease of use with high efficacy. External cooling methods are simple to use but slow in reducing core temperature. These techniques include the use of cooling blankets; application of ice packs to the groin, axillae, and neck; use of wet towels and fanning; and use of a cooling helmet [15]. In a recent study, intravenous infusion of 30 ml kg⁻¹ of crystalloid at 4 °C over 30 min reduced core temperature significantly and did not cause pulmonary oedema [26]. Cooling by peritoneal and pleural lavage is possible but not generally used [27]. Extracorporeal cooling methods are efficient [12] but too invasive for use in the prehospital environment or most emergency departments. An intravascular heat exchange device, which enables rapid cooling and precise temperature control, has recently become available.

Shivering during cooling leads to warming and will increase overall oxygen consumption. Shivering should be prevented by use of a neuromuscular blocker and sedation (as done in the two definitive trials). Careful monitoring of temperature is important during use of therapeutic hypothermia. The incidence of complications such as arrhythmias, infection, and coagulopathy is likely to increase if the core temperature falls considerably below 32 °C. Continuous monitoring of temperature can be accomplished by use of a bladder temperature probe or a pulmonary artery catheter if one is in situ. Other temperature-monitoring techniques, including intermittent tympanic temperature measurements, are less reliable.

5.4. Use of therapeutic hypothermia in children

There is currently insufficient evidence to make a recommendation on the use of therapeutic hypothermia in children resuscitated from

cardiac arrest. The European and Australian clinical trials excluded children and cardiac arrests of noncardiac aetiology (e.g. respiratory failure or shock), which are typical of those in children [18,19]. In the 1970s therapeutic hypothermia was used to reduce secondary brain injury in children with severe anoxic/ischaemic insults. The practice was abandoned in the 1980s after a retrospective study of near-drowning victims reported that children treated with hypothermia were at an increased risk for death, neutropenia, and sepsis compared with children treated without hypothermia [28,29]. Important limitations of this study were the limited sample size, use of historic controls, and use of a lower target temperature for a longer duration than recommended in contemporary protocols (30–33 °C for ≥ 36 h vs. 32–34 °C for 12–24 h).

Until additional paediatric data become available, clinicians should tailor therapy for individual patients based on their assessment of the risks and benefits of hypothermia. Risk-benefit assessment should take into account relevant data from laboratory models of asphyxial arrest [8,30–32], results from trials of adult cardiac arrest [18,19], and reports on the use of hypothermia in treatment of neonatal asphyxia [33–38]. The results of therapeutic hypothermia are generally favorable in laboratory models of hypoxic ischaemic injury to immature brains of various species. Preliminary data from clinical trials of perinatal asphyxia indicate that induced hypothermia is feasible and safe, but data on long-term neurological morbidity are not yet available [33–38]. It is difficult to extrapolate from these disparate sources of information, and thus there is no consensus yet for use of therapeutic hypothermia among clinicians who care for these critically ill children.

6. Summary: ILCOR recommendations

On the basis of the published evidence to date, the ILCOR ALS Task Force has made the following recommendations:

- Unconscious adult patients with spontaneous circulation after out-of-hospital cardiac arrest should be cooled to 32–34 °C for 12–24 h when the initial rhythm was VF.
- Such cooling may also be beneficial for other rhythms or in-hospital cardiac arrest.

Appendix A: 1998 AHA ECC levels of evidence summary

Level of evidence	Definitions
Level 1	One or more RCTs in which the lower limit of CI for treatment effect exceeds the minimal clinically important benefit
Level 2	One or more RCTs in which the lower limit of the CI for treatment effect overlaps the minimal clinically important benefit
Level 3	Prospective cohort of patients not randomized to an intervention; control or comparison group available
Level 4	Historic, nonrandomized cohort or case-control studies
Level 5	Case series: patients compiled in serial fashion; a control group is lacking
Level 6	An animal or mechanical model study; a level 6A study is well designed, shows a homogeneous pattern of results; a level 6B study has a less powerful design and demonstrates an equivocal or heterogeneous pattern of results
Level 7	Reasonable extrapolations from existing data; quasiexperimental designs; pathophysiological and nonquantitative reasoning
Level 8	Rational conjecture (common sense); common practices accepted before evidence-based guidelines

References

- [1] Benson DW, Williams GR, Spencer FC, Yates AJ. The use of hypothermia after cardiac arrest. *Anesth Analg* 1959;38:423–8.
- [2] Williams GR, Spencer FC. The clinical use of hypothermia following cardiac arrest. *Ann Surg* 1958;148:462–8.
- [3] Ravitch MM, Lane RR, Safar P, Steichen FM, Knowles P. Lightning strike: report of a case with recovery after cardiac massage and prolonged artificial respiration. *New Engl J Med* 1961;264:36–8.
- [4] Marion DW, Leonov Y, Ginsberg M, Katz LM, Kochanek PM, Lechleuthner A, Nemoto EM, Obrist W, Safar P, Sterz F, Tisherman SA, White RJ, Xiao F, Zar H. Resuscitative hypothermia. *Crit Care Med* 1996;24:81S–9S.
- [5] Horn M, Schlote W, Henrich HA. Global cerebral ischemia and subsequent selective hypothermia. *Acta Neuropathol (Berl)* 1991;81:443–9.
- [6] Sterz F, Safar P, Tisherman S, Radovsky A, Kuboyama K, Oku K. Mild hypothermic cardiopulmonary resuscitation improves outcome after prolonged cardiac arrest in dogs. *Crit Care Med* 1991;19:379–89.
- [7] D'Cruz BJ, Fertig KC, Filiano AJ, Hicks SD, DeFranco DB, Callaway CW. Hypothermic reperfusion after cardiac arrest augments brain-derived neurotrophic factor activation. *J Cereb Blood Flow Metab* 2002;22:848–51.

- [8] Hicks SD, DeFranco DB, Callaway CW. Hypothermia during reperfusion after asphyxial cardiac arrest improves functional recovery and selectively alters stress-induced protein expression. *J Cereb Blood Flow Metab* 2000;20:520–30.
- [9] Bernard SA, Jones BM, Horne MK. Clinical trial of induced hypothermia in comatose survivors of out-of-hospital cardiac arrest. *Ann Emerg Med* 1997;30:146–53.
- [10] Sanada T, Ueki M, Tokudome M, Okamura T, Nishiki S, Niimura A, Yabuki S, Watanabe Y. Recovery from out-of-hospital cardiac arrest after mild hypothermia: report of two cases. *Masui* 1998;47:742–5.
- [11] Yanagawa Y, Ishihara S, Norio H, Takino M, Kawakami M, Takasu A, Okamoto K, Kaneko N, Terei C, Okada Y. Preliminary clinical outcome study of mild resuscitative hypothermia after out-of-hospital cardiopulmonary arrest. *Resuscitation* 1998;39:61–6.
- [12] Nagao K, Hayashi N, Kanmatsuse K, Arima K, Ohtsuki J, Kikushima K, Watanabe I. Cardiopulmonary cerebral resuscitation using emergency cardiopulmonary bypass, coronary reperfusion therapy and mild hypothermia in patients with cardiac arrest outside the hospital. *J Am Coll Cardiol* 2000;36:776–83.
- [13] Zeiner A, Holzer M, Sterz F, Behringer W, Schorkhuber W, Mullner M, Frass M, Siostrzonek P, Ratheiser K, Kaff A, Laggner AN. Mild resuscitative hypothermia to improve neurological outcome after cardiac arrest: a clinical feasibility trial. Hypothermia After Cardiac Arrest (HACA) Study Group. *Stroke* 2000;31:86–94.
- [14] Felberg RA, Krieger DW, Chuang R, Persse DE, Burgin WS, Hickenbottom SL, Morgenstern LB, Rosales O, Grotta JC. Hypothermia after cardiac arrest: feasibility and safety of an external cooling protocol. *Circulation* 2001;104:1799–804.
- [15] Hachimi-Idrissi S, Corne L, Ebinger G, Michotte Y, Huyghens L. Mild hypothermia induced by a helmet device: a clinical feasibility study. *Resuscitation* 2001;51:275–81.
- [16] Callaway CW, Tadler SC, Katz LM, Lipinski CL, Brader E. Feasibility of external cranial cooling during out-of-hospital cardiac arrest. *Resuscitation* 2002;52:159–65.
- [17] The American Heart Association in Collaboration with the International Liaison Committee on Resuscitation (ILCOR). Guidelines 2000 for cardiopulmonary resuscitation and emergency cardiovascular care—an international consensus on science. *Resuscitation* 2000;46:1–447.
- [18] Bernard SA, Gray TW, Buist MD, Jones BM, Silvester W, Gutteridge G, Smith K. Treatment of comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *New Engl J Med* 2002;346:557–63.
- [19] The Hypothermia after Cardiac Arrest Study Group. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *New Engl J Med* 2002;346:549–56.
- [20] Holzer M. Therapeutic hypothermia after cardiac arrest [Letter]. *New Engl J Med* 2002;347:64.
- [21] Steen PA, Newberg L, Milde JH, Michenfelder JD. Hypothermia and barbiturates: individual and combined effects on canine cerebral oxygen consumption. *Anesthesiology* 1983;58:527–32.
- [22] Oku K, Sterz F, Safar P, Johnson D, Obrist W, Leonov Y, Kuboyama K, Tisherman SA, Stezoski SW. Mild hypothermia after cardiac arrest in dogs does not affect postarrest multifocal cerebral hypoperfusion. *Stroke* 1993;24:1590–7.
- [23] Colbourne F, Sutherland G, Corbett D. Postischemic hypothermia: a critical appraisal with implications for clinical treatment. *Mol Neurobiol* 1997;14:171–201.
- [24] Ginsberg MD, Sternau LL, Globus MY, Dietrich WD, Busto R. Therapeutic modulation of brain temperature: relevance to ischemic brain injury. *Cerebrovasc Brain Metab Rev* 1992;4:189–225.
- [25] Safar PJ, Kochanek PM. Therapeutic hypothermia after cardiac arrest. *New Engl J Med* 2002;346:612–3.
- [26] Bernard S, Buist M, Monteiro O, Smith K. Induced hypothermia using large volume, ice-cold intravenous fluid in comatose survivors of out-of-hospital cardiac arrest: a preliminary report. *Resuscitation* 2003;56:9–13.
- [27] Safar P, Behringer W, Böttiger BW, Sterz F. Cerebral resuscitation potentials for cardiac arrest. *Crit Care Med* 2002;30(Suppl. 4):S140–4.
- [28] Bohn DJ, Biggar WD, Smith CR, Conn AW, Barker GA. Influence of hypothermia, barbiturate therapy, and intracranial pressure monitoring on morbidity and mortality after near-drowning. *Crit Care Med* 1986;14:529–34.
- [29] Biggart MJ, Bohn DJ. Effect of hypothermia and cardiac arrest on outcome of near-drowning accidents in children. *J Pediatr* 1990;117:179–83.
- [30] Xiao F, Safar P, Radovsky A. Mild protective and resuscitative hypothermia for asphyxial cardiac arrest in rats. *Am J Emerg Med* 1998;16:17–25.
- [31] Hickey RW, Ferimer H, Alexander HL, Garman RH, Callaway CW, Hicks S, Safar P, Graham SH, Kochanek PM. Delayed, spontaneous hypothermia reduces neuronal damage after asphyxial cardiac arrest in rats. *Crit Care Med* 2000;28:3511–6.
- [32] Hachimi-Idrissi S, Corne L, Huyghens L. The effect of mild hypothermia and induced hypertension on long term survival rate and neurological outcome after asphyxial cardiac arrest in rats. *Resuscitation* 2001;49:73–82.
- [33] Gunn AJ, Gluckman PD, Gunn TR. Selective head cooling in newborn infants after perinatal asphyxia: a safety study. *Pediatrics* 1998;102:885–92.
- [34] Simbruner G, Haberl C, Harrison V, Linley L, Willeitner AE. Induced brain hypothermia in asphyxiated human newborn infants: a retrospective chart analysis of physiological and adverse effects. *Intensive Care Med* 1999;25:1111–7.
- [35] Azzopardi D, Robertson NJ, Cowan FM, Rutherford MA, Rampling M, Edwards AD. Pilot study of treatment with whole body hypothermia for neonatal encephalopathy. *Pediatrics* 2000;106:684–94.
- [36] Thoresen M, Whitelaw A. Cardiovascular changes during mild therapeutic hypothermia and rewarming in infants with hypoxic-ischemic encephalopathy. *Pediatrics* 2000;106:92–9.
- [37] Battin MR, Dezoete JA, Gunn TR, Gluckman PD, Gunn AJ. Neurodevelopmental outcome of infants treated with head cooling and mild hypothermia after perinatal asphyxia. *Pediatrics* 2001;107:480–4.
- [38] Compagnoni G, Pogliani L, Lista G, Castoldi F, Fontana P, Mosca F. Hypothermia reduces neurological damage in asphyxiated newborn infants. *Biol Neonate* 2002;82:222–7.